

Patterns of physical activity progression in patients with COPD

Statistical analysis full version

We compared the baseline characteristics of patients with follow-up vs those lost to follow-up by descriptive statistics and obtained p-values using mixed logistic regression models with random intercepts for study.

We identified cluster groups (physical activity patterns) using k-means (1), a hypothesis-free method that allowed grouping patients based on the baseline level, the final level and the change in daily step count. We used the Calinski-Harabasz stopping rule to decide the number of clusters (2). To characterize the patterns, we described physical activity and physical activity experience variables according to the cluster groups and compared baseline to follow-up values by paired t-tests. Because both studies used the same data collection methods main results are based on the pooled dataset and corrected for study.

To assess determinants of physical activity progression patterns, we first compared subjects' characteristics by physical activity patterns and obtained p-values from mixed logistic regression models with random intercepts for study and city area to account for possible heterogeneity in unmeasured characteristics related to study and city area. Then we built a multivariable multinomial regression model using the generalized linear latent and mixed model, with also random intercepts for study and city (3). Model building combined step-forward and backward algorithms, and determinants were included in the final model if: (i) they related to the outcome with a p-value <0.05 ; or (ii) they modified ($>10\%$ change in regression coefficient) the estimates of the remaining variables in the model (4). We tested goodness of fit of the final model.

We performed the following sensitivity analyses: (1) to investigate a possible difference between the two study samples, cluster analysis and description of resulting patterns

was performed separately for both samples; (2) to test whether the observed patterns were due to changes in wearing time, we tested the association between the change in daily step count and the change in wearing time overall and per pattern; (3) to rule out a relevant effect of pulmonary rehabilitation on the physical activity patterns we repeated the clustering after excluding patients included in pulmonary rehabilitation programs at baseline and/or during follow-up.

We estimated that the available sample size ($n=291$), fixed by the primary objectives of the original studies, was sufficient to identify physical activity patterns using cluster k-means, as our ratio of number of subjects to number of variables ($291/3 = 97$) was much higher than the 0.01 often used for the same analysis in other contexts (5,6).

Due to the small proportion of missing data (<2% of total data), we used a complete case strategy and reported missing data in the table footnotes.

All analyses were conducted using Stata/SE 14.2 (StataCorp, College Station, TX, USA).

Figure S1 Flow of participants through the study.




Table S1 Patient characteristics at baseline for all patients (Urban Training and PROactive study, n=412) and for patients with 12-month follow-up vs lost-to follow-up.

	All patients n = 412 (100%)	Follow-up n = 291 ^a (71%)	Lost-to follow-up n = 121 ^a (29%)	p- value ^b
Sociodemographic				
Age (years)	68±8	68±8	68±8	0.745
Sex (men)	316 (77)	237 (81)	79 (65)	0.001
Current smoker (yes)	77 (19)	52 (18)	25 (21)	0.508
Pack-years	58±41	58±41	60±41	0.684
Education, high school or higher	236 (57)	168 (58)	68 (56)	0.746
Interpersonal				
Living with a partner ^c	291 (71)	216 (74)	75 (63)	0.016
Active worker ^d	47 (11)	36 (12)	11 (9)	0.342
Grandparenting ^e	89 (43)	67 (45)	22 (39)	0.389
Dog walking ^e	26 (13)	20 (14)	6 (11)	0.566
Environmental				
Recruitment season				
Spring	55 (13)	35 (12)	20 (16)	
Summer	82 (20)	58 (20)	24 (20)	
Fall	218 (53)	154 (53)	64 (53)	0.161
Winter	57 (14)	44 (15)	13 (11)	
Urban vulnerability index (from 0 -lowest to 1 – highest) ^{e,f}	0.642±0.178	0.637±0.175	0.655±0.186	0.514
Clinical				
FEV ₁ (% predicted)	57.7±18.9	58.6±19.3	55.6±17.9	0.140
FEV ₁ /FVC ratio	0.51±0.13	0.51±0.13	0.51±0.13	0.699
Airflow limitation severity (post-bronchodilator FEV ₁)				
GOLD 1: Mild (FEV ₁ ≥ 80% predicted)	52 (13)	39 (13)	13 (11)	
GOLD 2: Moderate (50% ≤ FEV ₁ < 80% predicted)	207 (50)	147 (51)	60 (50)	
GOLD 3: Severe (30% ≤ FEV ₁ < 50% predicted)	125 (30)	88 (30)	37 (30)	0.259
GOLD 4: Very Severe (FEV ₁ <30% predicted)	28 (7)	17 (6)	11 (9)	
6MWD (meters)	461±109	477±103	421±111	<0.001
CAT score (0–40)	13.3±7.5	12.9±7.6	14.2±7.3	0.094
CCQ score (0-6)	1.59±0.98	1.55±0.98	1.69±0.98	0.172
C-PPAC amount score (0-100)	67.8±16.9	69.0±15.8	64.2±19.5	0.024
C-PPAC difficulty score (0-100)	77.9±14.9	78.4±14.5	76.3±16.0	0.269
C-PPAC total score (0-100)	72.8±13.6	73.7±12.8	70.3±15.4	0.044
mMRC score (0-4)	1.4±1.0	1.3±0.9	1.7±1.1	<0.001
Any COPD exacerbation with hospital admission in previous 12 months	49 (12)	34 (12)	15 (13)	0.781
BMI (kg/m ²)	27.7±5.2	27.6±4.6	28.1±6.3	0.306
FFMI (kg/m ²)	18.8±3.2	19.0±3.0	18.4±3.5	0.086
Cardiovascular disease ^g	240 (59)	176 (60)	64 (54)	0.212
Ischemic heart disease ^g	40 (10)	29 (10)	11 (9)	0.823
Diabetes mellitus ^g	73 (18)	51 (18)	22 (18)	0.817
LABA or LAMA, alone	56 (14)	41 (14)	15 (13)	0.686
Inhaled corticosteroid with LABA and/or LAMA	256 (63)	179 (62)	77 (65)	0.557
Pulmonary rehabilitation at baseline	25 (6)	15 (5)	10 (8)	0.233
Knowledge of baseline PA	24 (6)	19 (7)	5 (4)	0.348
Psychological				
Anxiety (HAD-A, 0-21)	5±4	5±4	6±4	0.117
Depression (HAD-D, 0-21)	4±3	4±3	4±4	0.210
Physical activity				
Step count (steps/day)	6415±3678	6720±3667	5682±3613	0.010
Time in moderate-to-vigorous physical activity (≥3 METs; min/day)	95.8±45.9	99.4±45.3	87.0±46.2	0.013
Intensity during walking (m/s ²)	1.84±0.31	1.86±0.31	1.80±0.30	0.050
Sedentary time (h/day)	10.53±1.93	10.53±1.94	10.52±1.92	0.961

Notes: Data are presented as n (%), mean±SD.

^aSome variables have missing values, as follows. Follow-up: 1 in education, 1 in living with a partner, 1 in CAT total, 1 in CCQ score, 31 in C-PPAC scores, 1 in any COPD exacerbation with hospital admission in previous 12 months, 26 in FFMI, 3 in LABA or LAMA, alone, 3 in inhaled corticosteroid with LABA and/or LAMA, 3 in HAD anxiety and depression; Lost-to follow-up: 1 in living with a partner, 1 in 6MWD, 33 in C-PPAC scores, 3 in any COPD exacerbation with hospital admission in previous 12 months, 5 in FFMI, 2 in ICD10 codes: I00 to I99 for Cardiovascular diseases; I20 to I25 for Ischemic heart disease, E14 for Diabetes mellitus, 3 in LABA or LAMA, alone, 3 in inhaled corticosteroid with LABA and/or LAMA, 1 in HAD depression.

^bp-value from mixed logistic regression models with random effects for study (Urban Training and PROactive), due to small numbers random effects for city area were not applied.

^cmarital status: living with a partner vs single, widowed or divorced.

^dworking status: active worker (working full-time or part-time) vs. unemployed, housework or retired.

^eonly available for Urban Training.

^fThe urban vulnerability index is a measure of socioeconomic status at the census tract level that combines demographic, economic, residential and subjective indicators, and ranges from lowest [0] to highest [1] level of neighborhood vulnerability.

^gICD10 codes: I00 to I99 for cardiovascular diseases; I20 to I25 for ischemic heart disease, E14 for diabetes mellitus.

Abbreviations: FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; GOLD: Global Initiative for Chronic Obstructive Lung Disease; 6MWD: 6-min walking distance; CAT: COPD Assessment Test; CCQ: Clinical COPD Questionnaire; C-PPAC: Clinical visit—PROactive Physical Activity in COPD (higher numbers indicate a better score); mMRC: modified Medical Research Council; BMI: body mass index; FFMI: fat free mass index; LABA: long-acting beta₂-agonists; LAMA: long-acting anti-muscarinics; HAD-A: Hospital Anxiety and Depression scale - Anxiety; HAD-D: Hospital Anxiety and Depression scale – Depression; MET: metabolic equivalent of task.

Table S2 Physical activity and physical activity experience variables at baseline and at 12-month follow-up, overall and by PA progression pattern (*Inactive, Active Improvers and Active Decliners*).

	<i>All</i>				<i>Inactive</i>				<i>Active Improvers</i>				<i>Active Decliners</i>			
	n = 291 ^a				n = 173 ^a (59%)				n = 49 ^a (17%)				n = 69 ^a (24%)			
	Baseline	Follow-up	Change	p-value ^b	Baseline	Follow-up	Change	p-value ^b	Baseline	Follow-up	Change	p-value ^b	Baseline	Follow-up	Change	p-value ^b
Step count (steps/day)	6720 ±3667	6474 ±3772	-246 ±2420	0.084	4621 ±1757	4134 ±1817	-487 ±1201	<0.001	7727 ±3275	11105 ±3330	3378 ±2203	<0.001	11267 ±3009	9051 ±2897	-2217 ±2085	<0.001
Time in MVPA (≥3 METs; min/day)	99.4 ±45.3	96.1 ±47.5	-3.3 ±29.1	0.052	74.5 ±25.9	68.2 ±26.6	-6.3 ±16.7	<0.001	114.8 ±47.2	152.7 ±47.1	37.9 ±26.8	<0.001	150.9 ±32.3	125.8 ±33.0	-25.0 ±26.0	<0.001
Intensity during walking (m/s ²)	1.86 ±0.31	1.82 ±0.31	-0.04 ±0.19	<0.001	1.78 ±0.27	1.73 ±0.27	-0.05 ±0.19	<0.001	1.90 ±0.34	1.95 ±0.32	0.05 ±0.17	0.062	2.03 ±0.30	1.95 ±0.30	-0.08 ±0.17	<0.001
Sedentary time (h/day)	10.53 ±1.94	10.46 ±1.98	-0.08 ±1.82	0.463	11.03 ±2.00	10.98 ±2.15	-0.05 ±2.10	0.744	10.29 ±1.73	9.59 ±1.54	-0.70 ±1.25	<0.001	9.46 ±1.39	9.76 ±1.29	0.30 ±1.17	0.037
C-PPAC amount score (0-100)	67.6 ±15.9	66.7 ±16.0	-0.9 ±12.7	0.305	61.5 ±14.2	60.3 ±14.7	-1.2 ±12.4	0.255	73.6 ±12.0	82.3 ±10.3	8.7 ±11.2	<0.001	83.6 ±10.3	77.1 ±9.4	-6.5 ±11.0	<0.001
C-PPAC difficulty score (0-100)	77.2 ±14.3	77.7 ±13.6	0.4 ±10.6	0.547	74.4 ±14.2	74.8 ±13.0	0.4 ±10.0	0.673	80.3 ±13.8	84.5 ±11.3	4.3 ±11.3	0.054	84.4 ±12.5	82.5 ±14.0	-1.9 ±11.5	0.295
C-PPAC total score (0-100)	72.4 ±12.6	72.2 ±12.6	-0.2 ±9.2	0.717	68.0 ±11.7	67.5 ±11.5	-0.4 ±8.5	0.562	76.9 ±9.5	83.4 ±8.8	6.5 ±8.0	<0.001	84.0 ±8.1	79.8 ±9.6	-4.2 ±9.7	0.008
Wearing time (h/day)	14.73 ±1.56	14.52 ±1.63	-0.21 ±1.67	0.035	14.68 ±1.72	14.35 ±1.82	-0.33 ±2.01	0.034	14.77 ±1.58	15.02 ±1.59	0.25 ±0.77	0.028	14.82 ±1.10	14.59 ±0.93	-0.23 ±1.03	0.064

Notes: Data are presented as mean±SD. For C-PPAC variables means and p-values are reported for patients with data at baseline and follow-up.

^aC-PPAC variables have 87 missing values: 38 in *Inactive*, 21 in *Active Improvers*, and 28 in *Active Decliners*.

^bpaired t-test.

Abbreviations: MVPA: moderate-to-vigorous physical activity; MET: metabolic equivalent of task; C-PPAC: Clinical visit—PROactive Physical Activity in COPD (higher numbers indicate a better score).

Table S3 Step count (mean steps/day) at baseline and at 12-month follow-up as well as selected variables at baseline, by cluster groups (physical activity progression patterns) identified by k-means, performed separately for the Urban Training and the PROactive study.

	<i>Cluster 1</i>					<i>Cluster 2</i>					<i>Cluster 3</i>				
	n (row%)	Baseline	Follow-up	Change	p-value ^a	n (row%)	Baseline	Follow-up	Change	p-value ^a	n (row%)	Baseline	Follow-up	Change	p-value ^a
Urban Training															
Step count (steps/day)	79 (54%)	6028 ±2176	5125 ±2045	-903 ±1428	<0.001	36 (24%)	6722 ±2491	10435 ±2293	3713 ±2288	<0.001	33 (22%)	13144 ±3340	11442 ±3606	-1702 ±2754	0.001
Age (years)		69±8					71±9					67±7			
Sex (men)		70 (89)					32 (89)					28 (85)			
FEV ₁ (% predicted)		55.2±18.2					62.6±15.5					60.5±17.3			
6MWD (meters)		485±87					519±83					517±68			
mMRC score (0-4)		1.2±0.9					1.1±0.8					0.8±0.7			
PROactive															
Step count (steps/day)	118 (83%)	4431 ±1735	4097 ±1891	-334 ±1098	0.001	6 (4%)	11339 ±1239	14340 ±2593	3001 ±1716	0.008	19 (13%)	11188 ±1992	8222 ±2685	-2966 ±2166	<0.001
Age (years)		68±8					65±9					63±7			
Sex (men)		89 (75)					4 (67)					14 (74)			
FEV ₁ (% predicted)		56.8±20.7					68.6±12.1					69.8±21.7			
6MWD (meters)		435±109					520±130					541±110			
mMRC score (0-4)		1.6±1.0					1.0±0.6					1.0±0.8			

Notes: Data are presented as mean±SD.

^apaired t-test.

Abbreviations: FEV₁: forced expiratory volume in 1 second; 6MWD: 6-min walking distance; mMRC: modified Medical Research Council.

Table S4 Correlation between the change in daily step count and the change in wearing time, overall and by PA progression pattern (*Inactive*, *Active Improvers* and *Active Decliners*).

	n (%)	Pearson correlation coefficient	p-value
<i>All patients</i>	291 (100%)	0.090	0.124
<i>Inactive</i>	173 (59%)	0.002	0.981
<i>Active Improvers</i>	49 (17%)	-0.097	0.508
<i>Active Decliners</i>	69 (24%)	0.162	0.184

Table S5 Step count (mean steps/day) at baseline and at 12-month follow-up, by cluster groups (physical activity progression patterns) identified by k-means, performed separately for all patients (n=291) and excluding patients with rehabilitation (n=270).

	<i>Inactive</i>					<i>Active Improvers</i>					<i>Active Decliners</i>				
	n (row%)	Baseline	Follow-up	Change	p-value ^b	n (row%)	Baseline	Follow-up	Change	p-value ^b	n (row%)	Baseline	Follow-up	Change	p-value ^b
Step count (steps/day), all patients	173 (59%)	4621 ±1757	4134 ±1817	-487 ±1201	<0.001	49 (17%)	7727 ±3275	11105 ±3330	3378 ±2203	<0.001	69 (24%)	11267 ±3009	9051 ±2897	-2217 ±2085	<0.001
Step count (steps/day), patients with rehabilitation excluded ^a	154 (57%)	4544 ±1763	4003 ±1753	-541 ±1209	<0.001	50 (19%)	7516 ±3187	10730 ±3345	3213 ±2266	<0.001	66 (24%)	11206 ±3033	9078 ±2928	-2128 ±1943	<0.001

Notes: Data are presented as mean±SD.

^apatients in pulmonary rehabilitation at baseline and/or follow-up were excluded (n=21).

^bpaired t-test

References:

1. Genolini, Christophe Falissard B. KmL: k-means for longitudinal data. *Comput Stat* 2010;25:317–328.
2. Calinski T, Harabasz J. A dendrite method for cluster analysis. *Commun Stat - Theory Methods* 1974;3:1–27.
3. Rabe-Hesketh S, Skrondal A, Pickles A. Maximum likelihood estimation of limited and discrete dependent variable models with nested random effects. *J Econom* 2005;128:301–23.
4. Hosmer DW, Lemeshow S, Sturdivant RX. *Applied logistic regression*. 3rd ed. Hoboken, New Jersey: John Wiley & Sons; 2013.
5. Wang Y, Miller DJ, Clarke R. Approaches to working in high-dimensional data spaces: Gene expression microarrays. *Br J Cancer* 2008;98:1023–8.
6. Garge NR, Page GP, Sprague AP, Gorman BS, Allison DB. Reproducible clusters from microarray research: Whither? *BMC Bioinformatics* 2005;6:S10.