



Supplementary material

The ERICE-Score: the New Native Cardiovascular Score for the low-risk and Aged Mediterranean Population of Spain

Rafael Gabriel ^{a,b,*}, Carlos Brotons ^{b,c}, M. José Tormo ^{d-f}, Antonio Segura ^g,
Fernando Rigo ^{h,i}, Roberto Elosua ^{e,j}, Julio A. Carbayo ^k, Diana Gavrila ^{d-f},
Irene Moral ^c, Jaakko Tuomilehto ^{a,b,l} y Javier Muñoz ^{a,b,m}; on behalf of the ERICE
Study Group [◇]

^a*Unidad de Epidemiología Clínica, Instituto IdiPAZ, Hospital Universitario La Paz, Universidad Autónoma de Madrid, Madrid, Spain*

^b*Red de Investigación Cardiovascular (RIC), Spain*

^c*Unidad de Investigación, EAP Sardenya-IIB Sant Pau, Barcelona, Spain*

^d*Servicio de Epidemiología, Consejería de Sanidad de Murcia, Murcia, Spain*

^e*CIBER Epidemiología y Salud Pública, CIBEResp, Spain*

^f*Departamento de Ciencias Sociosanitarias, Universidad de Murcia, Murcia, Spain*

^g*Instituto de Ciencias de la Salud de Castilla-La Mancha, Talavera de la Reina, Toledo, Spain*

^h*Grupo CORSAIB, IB-Salut, Palma de Mallorca, Islas Baleares, Spain*

ⁱ*Red de Investigación en Actividades Preventivas y Promoción de la Salud (redIAPP), Spain*

^j*Epidemiología Cardiovascular y Genética, IMIM (Instituto de Investigación Hospital del Mar), Barcelona, Spain*

^k*Unidad de Lípidos, Clínica Nuestra Señora del Rosario, Grupo de Enfermedades Vasculares de Albacete (GEVA), Albacete, Spain*

^l*University of Helsinki, Helsinki, Finland*

^m*Instituto Universitario de Ciencias de la Salud, Universidad de A Coruña, A Coruña, Spain*

ERICE Study Group members

Hospital Universitario La Paz and Hospital Universitario de La Princesa, Madrid (Coordinating Center): Rafael Gabriel, Francisco Rodríguez-Salvanés, Guillermo Fernández, Margarita Alonso, Carmen Suárez, Río Aguilar, Blanca Novella, Saturio Vega, Ana Ortega, Sagrario Susi, Rosario Madero, Eva Martínez Renedo, Jaakko Tuomilehto.

Instituto Universitario de Ciencias de la Salud Universidad de A Coruña and Complejo Hospitalario Universitario de A Coruña, La Coruña: Javier Muñiz, Isidro López, Teresa R. Pérez-Castro, Pilar Rodríguez, María L. Zapata, X. Antonio Méndez.

Hospital de Vic, Barcelona and EAP Sardenya-IIB Sant Pau, Barcelona: Carlos Brotons, Pere Roura, Alfonso Leyva, Emilia Chirveches, Rosa Salla, Irene Moral

Instituto Municipal de Investigaciones Médicas, Barcelona: Jaume Marrugat, Roberto Elosua, Izabella Rolhfs, Helena Martí

Grupo de Enfermedades Vasculares de Albacete [GEVA], Albacete: Juan A. División, Miguel Artigao, Esperanza Martínez, Carlos Sanchis, Julio A. Carbayo, Lucinio Carrión, Javier Massó, Enrique López de Coca, Francisco García Gosálvez, Beatriz Rodríguez-Paños, David Caldevilla, Juan López-Abril, Rosalina Martínez, Eva Argandoña, Alfonso Campayo.

Instituto de Ciencias de la Salud, Talavera de la Reina, Toledo: Antonio Segura, José María García, Marta Rodríguez, Carmen García, Enrique Almar, Antonio Mateos, Gema Rius, Luis Rodríguez, Eduardo Castellanos, Juan E. Alcalá, Catherine Graupner, Pedro Hernández, Luis F. Pajín, Tomás Cantón, José Moreu, Carolina Maicas, Ana M. García, Gema Vega

Consejería de Sanidad, Murcia: María J. Tormo, Diana Gavrila, José A. Melgarejo, José García, Miguel Rodríguez, C Navarro.

Dirección de Atención Primaria Mallorca, Grupo CORSAIB, RedIAPP: Bartolomé Seguí, Fernando Rigo, Guillem Frontera, Tomás Rodríguez, Francisco J. Sánchez, Isabel Borràs, Joan Llovera, Alfredo Sebastián, Manuel Rullan, Pablo Hermoso.

CALCULATION OF 10-YEAR RISK ESTIMATES FOR CARDIOVASCULAR DISEASE

Step 1

Calculate the underlying risk factors for CVD events (fatal and nonfatal) using a bivariate analysis. Of the originally entered variables that were selected based on theoretical information, the nonsignificant ones (estimated $p > .05$) were dropped from consideration.

Step 2

Cox proportional-hazard regressions analyzing the effect of the selected risk factors on survival were carried out. Cox models were adjusted by study source. Separate hazard functions for males and females were performed. The risk factors were introduced as stratified variables because analysis of heterogeneity was positive.

Cox models stratified by age ≤ 70 years or > 70 years were tested and C-statistic showed a good discriminative power (C statistic > 0.80). However, the model by event and sex reported a better estimation of the risk probabilities in the score calculation (C statistic > 0.78). Omnibus test was used to validate the model ($P < .05$). The C statistic, which is equivalent to areas under receiver operator characteristic (ROC) curves, is a measure of discrimination that estimates the ability of a predictive model to differentiate individuals who experience an event from those who did not.

Step 3

Weighted sums for men and women are calculated. The sums ($\sum \beta_i x_i$) are obtained replacing β_i by the β -coefficients obtained in the Cox regression models for each evaluated risk factor (coefficients in tables 3 for men and in table 4 for women), and replacing x_i by "1" when the risk factor is present or "0" when it is not.

Step 4

Survival is defined as the proportion of patients living for a certain amount of time (10 years).

Probability of an event happening within the next 10 years was calculated as follows:

$$P_{X_i(t)} = 1 - S_{x_i(t)} = 1 - e^{-[H_{0(t)} \cdot e^{\sum(\beta_i x_i) - \sum(\beta_i x_m)}]} = 1 - e^{-[-\ln(S_{0(t)}) \cdot e^{\sum(\beta_i x_i) - \sum(\beta_i x_m)}]} = 1 - S_{0(t)} e^{\sum(\beta_i x_i) - \sum(\beta_i x_m)}$$

Where $e^{\sum(\beta_i x_i)}$ is the exponential of the sum of beta coefficients from the risk factors that are present in the person being evaluated, $e^{\sum(\beta_i x_m)}$ is the exponential of the sum of beta coefficients from the risk factors that are present in the average population and $S_{0(t)}$ is the baseline probability of being event-free during a time (t).

Using Kaplan-Meier methods, 10-year survival probability of 88.14% was estimated for men and 91.15% for women.

Table 1 Supplementary Material

Distribution of Cardiovascular Risk Factors in the ERICE Study by Sex

		Men		Women		Total	
		N	%	n	%	n	%
		5413	45.9	6387	54.1		
Age	<40	963	17.8	972	15.2	1935	16.4
	40-49	1006	18.6	1233	19.3	2239	19
	50-59	825	15.2	937	14.7	1762	14.9
	60-69	1345	24.8	1520	23.8	2865	24.3
	70-79	965	17.8	1253	19.6	2218	18.8
	≥80	309	5.7	472	7.4	781	6.6
Systolic blood pressure	< 140	3258	60.2	3671	57.5	6929	58.7
	140-160	1469	27.1	1748	27.4	3217	27.3
	160-180	542	10	727	11.4	1269	10.8
	≥180	144	2.7	241	3.8	385	3.3
Cholesterol	< 200	2065	38.1	2290	35.9	4355	36.9
	200-249	2368	43.7	2764	43.3	5132	43.5
	250-299	837	15.5	1121	17.6	1958	16.6
	≥300	143	2.6	212	3.3	355	3
Diabetics		479	8.8	538	8.4	1017	8.6
Smoking		3328	61.9	964	15.2	4292	36.4

The value obtained is multiplied by 100 to estimate the percentage of risk that individuals with the characteristics described (age, systolic blood pressure, smoking, total cholesterol, diabetes) will develop a cardiovascular event within the next 10 years.

Table 2 Supplementary Material

Bivariate Model for a First Fatal or Nonfatal Cardiovascular Event

		Men				Women			
		Sig.	Exp(B)	95%CI		Sig.	Exp(B)	95%CI	
				Lower	Upper			Lower	Upper
Age		0.000	1.072	1.064	1.080	0.000	1.096	1.087	1.105
Smoking	No		1.000				1.000		
	Yes	0.052	1.205	0.998	1.453	0.000	0.440	0.344	0.562
Diabetes	No		1.000				1.000		
	Yes	0.000	1.949	1.532	2.480	0.000	2.472	1.945	3.141
Systolic blood pressure		0.000	1.024	1.020	1.028	0.000	1.023	1.019	1.027
Cholesterol		0.639	1.000	0.999	1.002	0.000	1.004	1.002	1.006
Body mass index		0.768	1.003	0.982	1.025	0.031	1.019	1.002	1.036
N		5260				6187			