



Supplementary material

**Awareness of Genetic Coronary Risk Score Improves Blood Pressure Control
in Hypertensive Patients**

SUPPLEMENTARY MATERIAL

Table of the supplementary material.

Ambulatory Blood Pressure Values at Baseline and Changes at the End Study

Blood pressure, mmHg		Genetic risk score group (n = 31)	Control group (n = 36)	<i>P</i> *
Systolic				
24 h	Baseline	140.0 (138.03 to 51.07)	139.62 (130.92 to 48.32)	.866
	Change	-16.31(-26.03 to -6.59)	-14,52(-23.5 to -5.54)	.442
Daytime	Baseline	145.1 (156.3 to 133.9)	145.5 (154.5 to 136.5)	.882
	Change	-18.43 (-28.72 to -8.14)	-15.67 (-25.96 to -5.38)	.290
Nighttime	Baseline	127.3 (140.3 to 114.3)	126.7 (139.1 to 114.3)	.846
	Change	-12.15 (-23.92 to -0.38)	-11.58 (-22.44 to 1.02)	.676
Diastolic				
24 h	Baseline	88.84 (96.14 to 81.54)	87.41 (94.51 to 80.31)	.425
	Change	-10.96 (-18.88 to -3.04)	-8.50 (-15.62 to 1.38)	.118
Daytime	Baseline	93.4 (100.5 to 86.3)	92.3 (99.4 to 85.2)	.552
	Change	-11.93 (-18.53 to -5.33)	-9.11 (-17.0 to -1.21)	.123
Nighttime	Baseline	78.3 (87.5 to 61.1)	76.4 (83.3 to 69.5)	.347
	Change	-8.28 (-16.3 to -0.26)	-7.88 (-15.29 to 0.47)	.645

Data are expressed as mean (95% confidence interval). Changes for which the 95% confidence interval does not include 0 are significantly different from the baseline. * *P* value for comparisons among groups with 1-way ANOVA.