

SUPPLEMENTARY MATERIAL

Table 1 of the supplementary material

Recalculated Predictive Model After Exclusion of Patients Who Died During Hospital Admission

	OR (95%CI)	<i>P</i>
non-Shockable rhythm	11.79 (3.16-18.63)	<.01
Age (one additional year)	1.05 (1.01-1.12)	.01
Lactate level (one additional mmol/L)	1.23 (1.09-1.43)	.04
Time Elapsed to return of spontaneous circulation	1.02 (1.00-1.06)	.03
Diabetes	3.65 (1.56-12.54)	.02

95%CI, confidence interval; OR, adjusted odds ratio.

Table 2 of the supplementary material


Recalculated Predictive Model After Exclusion of Patients With Noncardiac Causes of out-of-Hospital Cardiac Arrest

	OR (95%CI)	<i>P</i>
non-Shockable rhythm	43.01 (39.43-47.08)	<.01
Age (1 additional year)	1.05 (1.01-1.09)	.03
Lactate level (one additional mmol/L)	1.55 (1.20-2.00)	<.01
Time Elapsed to return of spontaneous circulation	1.05 (1.00-1.09)	.04
Diabetes	6.57 (1.55-12.93)	.01

95%CI, confidence interval; OR, adjusted odds ratio.

Figure of the supplementary material

Receiver operator characteristic curve for predictive model applied to in hospital survivors (n = 115).



95%CI, confidence interval; ROC, receiver operator characteristic.