

Prognostic value of lung ultrasound in chronic stable ambulatory heart failure patients

SUPPLEMENTARY DATA

Table 1 of the supplementary data

Univariable Cox regression analysis for the composite endpoint and for all-cause death

| | Composite endpoint ^a | | | All-cause death | | |
|--|---------------------------------|-----------|-------|-----------------|-----------|-------|
| | HR | 95% CI | P | HR | 95% CI | P |
| <i>Total B-lines sum</i> | 1.05 | 1.03–1.08 | <.001 | 1.06 | 1.04–1.09 | <.001 |
| <i>Age, y</i> | 1.05 | 1.04–1.08 | <.001 | 1.07 | 1.05–1.09 | <.001 |
| <i>Female sex</i> | 1.12 | 0.80–1.58 | .52 | 1.15 | 0.77–1.73 | .49 |
| <i>Ischemic etiology</i> | 1.83 | 1.33–2.50 | <.001 | 1.97 | 1.35–2.87 | <.001 |
| <i>HF duration, y^b</i> | 1.27 | 1.06–1.54 | .01 | 1.20 | 0.97–1.50 | .10 |
| <i>NYHA class</i> | 2.83 | 2.12–3.78 | <.001 | 2.99 | 2.12–4.21 | <.001 |
| <i>LVEF, %</i> | 0.98 | 0.97–0.99 | .002 | 0.98 | 0.97–0.99 | .02 |
| <i>Diabetes</i> | 1.75 | 1.28–2.39 | <.001 | 1.67 | 1.15–2.43 | .007 |
| <i>Hypertension</i> | 1.75 | 1.25–2.46 | .001 | 1.79 | 1.19–2.69 | .005 |
| <i>COPD</i> | 1.31 | 0.83–2.08 | .25 | 1.28 | 0.74–2.21 | .37 |
| <i>Atrial fibrillation/flutter</i> | 1.72 | 1.21–2.43 | .002 | 1.57 | 1.04–2.39 | .03 |
| <i>Anemia^c</i> | 2.17 | 1.58–2.98 | <.001 | 2.82 | 1.94–4.10 | <.001 |
| <i>Renal insufficiency^d</i> | 2.87 | 2.05–4.02 | <.001 | 2.66 | 1.78–3.97 | <.001 |
| <i>BMI, kg/m²</i> | 0.97 | 0.94–1.00 | .09 | 0.95 | 0.92–1.00 | .03 |
| <i>NT-proBNP, ng/L^e</i> | 2.10 | 1.80–2.45 | <.001 | 2.25 | 1.87–2.72 | <.001 |
| <i>Treatments</i> | | | | | | |
| <i>ACEI or ARB</i> | 0.37 | 0.26–0.51 | <.001 | 0.38 | 0.26–0.56 | <.001 |
| <i>Beta-blocker</i> | 0.68 | 0.42–1.16 | .13 | 0.51 | 0.30–0.86 | .01 |

| | | | | | | |
|----------------------|------|-----------|-------|------|-----------|-------|
| MRA | 1.20 | 0.87–1.66 | .26 | 1.20 | 1.15–1.68 | 0.79 |
| Sacubitril/valsartan | 0.94 | 0.30–2.96 | .92 | 0.43 | 0.06–3.06 | .40 |
| Loop diuretic | 2.22 | 1.49–3.31 | <.001 | 2.04 | 1.28–3.26 | .003 |
| Digoxine | 1.30 | 0.89–1.89 | .18 | 1.00 | 0.62–1.61 | .99 |
| Ivabradine | 0.84 | 0.53–1.31 | .43 | 0.86 | 0.51–1.46 | .58 |
| Hydralazine | 2.79 | 1.96–3.96 | <.001 | 3.40 | 2.28–5.06 | <.001 |
| Nitrates | 2.53 | 1.84–3.49 | <.001 | 2.58 | 1.77–3.77 | <.001 |
| CRT | 1.14 | 0.76–1.70 | .54 | 1.37 | 0.86–2.16 | .18 |
| IDC | 0.93 | 0.64–1.34 | .69 | 0.83 | 0.53–1.31 | .42 |

ACEI, angiotensin converting enzyme inhibitor; ARB, angiotensin II receptor blocker; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CRT, cardiac resynchronization therapy; HF, heart failure; ICD, implantable cardiac defibrillator; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; NYHA, New York Heart Association; NT-proBNP, N-terminal pro-brain natriuretic peptide.

^a Composite of all-cause death or HF hospitalization.

^b Log-transformed and in months.

^c According to WHO criteria (< 13 g/dL in men and < 12 g/dL in women).

^d Estimated glomerular filtration rate (Chronic Kidney Disease Epidemiology Collaboration equation) <60 mL/min per 1.73 m².

^e Log-transformed and for 1 standard deviation.

Table 2 of the supplementary data

Study design, LUS-specific components and results

| | |
|------------------------------|--|
| <i>Patients</i> | <ul style="list-style-type: none"> • Heart failure clinic • Ambulatory stable HF patients • Excluded patients: pulmonary fibrosis or radiological diffuse pleural fibrosis |
| <i>LUS image adquisition</i> | <ul style="list-style-type: none"> • Pocket device (V-scan simple model with a single sector probe, General Electric) • Phased array transducer, perpendicular to the ribs and an imaging depth of 14 cm • Patient in a supine position • Protocol: 8 areas |
| <i>Lus image analysis</i> | <ul style="list-style-type: none"> • Off-line analysis • Sonographer and reader blinded to clinical data, NT-proBNP and echocardiogram • Number of B-lines in each thoracic area • The sum of B-lines across all lung areas and the quartiles of such addition were used for the analyses |
| <i>Data analyses</i> | <ul style="list-style-type: none"> • Main clinical outcomes: composite endpoint of all-cause death or HF hospitalization and mortality from any cause • Follow-up: mean 31 ± 7 mo • Mean number of B-lines: 5 ± 6 • Q1, 0; Q2, 1-3; Q3, 4-7; Q4, ≥ 8 • Survival curves: having ≥ 8 B-lines (Q4) doubled the risk of experiencing the composite endpoint ($P < .001$) and increased by 2.6-fold the risk of death from any cause ($P < .001$) • Multivariable analysis: sum of B-lines across all lung areas remained as an independent prognostic factor of the composite endpoint and all-cause death, independently of the inclusion of NT-proBNP in the model |

LUS, lung ultrasound; NT-proBNP, N-terminal pro-brain natriuretic peptide.

FIGURES OF THE SUPPLEMENTARY DATA




Figure 1 of the supplementary data. Distribution of sum of B-lines across all lung areas per patient in the total cohort. Around one fourth of the patients had 0 B-lines.




Figure 2 of the supplementary data. Scatter-plot representing the sum of B-lines (X axis) and logNTproBNP (Y axis). A statistically significant but rather modest correlation was found.




Figure 3 of the supplementary data. Calibration plots for the composite endpoint in predictive models with and without NT-proBNP (A and B) and for all-cause death in both models (C and D, respectively). Calibration was assessed by plotting the predicted probability of the clinical endpoint against the observed frequency.