Figure 1 ESM. Study selection for this systematic review of the APP in COVID-19 subjects.




Figure 2 ESM. Sensitivity analysis: exclusion of study with large effect size.

	Awake prone positioning		Usual Care		Risk Ratio		Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% C	M-H, Fixed, 95% CI
Gad/Egypt	3	15	3	15	11.1%	1.00 [0.24, 4.18]	
Jayakumar/India	4	30	4	30	14.7%	1.00 [0.28, 3.63]	
Johnson/USA	2	15	1	15	3.7%	2.00 [0.20, 19.78]	
Rosén/Sweden	12	36	13	39	46.0%	1.00 [0.53, 1.90]	
Sryma/India	2	30	5	15	24.6%	0.20 [0.04, 0.91]	· · · · · · · · · · · · · · · · · · ·
Taylor/USA	o	27	0	13		Not estimable	
Total (95% CI)		153		127	100.0%	0.84 [0.52, 1.35]	•
Total events	23		26				
Heterogeneity: Chi ² =	4.39, df = 4 (P = 0.36)	; I2 = 9%					201 21 10 10
Test for overall effect: Z = 0.72 (P = 0.47)							0.01 0.1 1 10 100 Awake prone positioning Usual Care